

BEXSERO DOSING SCHEDULE

The doctor, pharmacist or nurse will inject BEXSERO into the arm or leg muscle, depending on the age of the patient.

RECOMMENDED VACCINATION SCHEDULE:

Age group at time of first dose	Primary immunization		Booster dose (0.5 mL)
	Primary doses <i>Each dose is 0.5 mL</i>	Interval between primary doses	
Infants (Option 1) 2 through 5 months of age	2+1 dosing schedule (Option 1)		1 dose Required in the 2nd year of life after an interval of ≥6 months from the last injection of the initial course.
	2	≥2 months	
Infants (Option 2) 2 through 5 months of age	3+1 dosing schedule (Option 2)		1 dose Required in the 2nd year of life after an interval of ≥6 months from the last injection of the initial course.
	3	≥1 month	
Infants 6 through 11 months of age	2	≥2 months	1 dose Required in the 2nd year of life, after an interval of ≥2 months from the last dose.
Children 12 months through 23 months	2	≥2 months	1 dose Required after an interval of 12 to 23 months from the second injection.
Children, adolescents and adults 2 through 25 years of age	2	≥1 month	1 dose Considered in individuals at continuous risk of exposure to meningococcal disease.

Vaccination against MenB* is not part of your child's routine vaccination schedule. Help protect your child against MenB* with BEXSERO.

Please refer to the package insert for complete dosing information.

* Meningococcal disease caused by *Neisseria meningitidis* group B strains.

Date of initial dose: _____ / ____ / ____